The background of the slide is a dark blue, textured surface resembling a spiral-bound notebook. At the top, there is a black horizontal band with a series of white, spiral-shaped rings, mimicking the binding of a notebook. The main title is centered on the page in a large, white, sans-serif font.

# Hubble Residuals and Host Galaxy Metallicities in an 'Unbiased' Sample

Chris D'Andrea  
University of Pennsylvania  
October 25th, 2010  
SDSS-II SNS Collaboration Meeting




# Motivation


2

- ◆ Dependence of SN Ia stretch, rates on Host Type have been shown (Gallagher et al. 2005; Sullivan et al. 2006)
- ◆ More recently, overluminosity *after* LC corrections has also been shown to depend on Host properties (Gallagher et al. 2008; Kelly et al. 2010; Sullivan et al. 2010; Lampeitl et al. 2010)

Kelly et al. (2010)



Lampeitl et al. (2010)






# Motivation


3

- ◆ SNe Ia overall luminosity depends on progenitor metallicity, with higher  $Z$  producing less  $^{56}\text{Ni}$  and a less luminous event (Timmes et al. 2003)
- ◆ For a given absolute magnitude, a SN Ia with a higher progenitor metallicity will have a *narrower* LC (Kasen, Ropke, & Woosley 2009)
- ◆ Believed that the Host Mass - Hubble Residual correlation is due to progenitor metallicity
- ◆ *Has not been directly measured!*

Sullivan et al. (2010)



Tremonti et al. (2010)






# SDSS-II SNS sample

4

- ◆ 'Completeness' - refers to SNe Ia detection efficiency @  $z < 0.15$  (Dilday et al. 2008)
- ◆ Complete sample for this study would be limited to those SNe with Distance Modulus from Light Curve fit
- ◆ Galaxy-only spectra for 88% of hosts
- ◆ Spectra from SDSS-I; BOSS; NTT; APO; GEMINI
- ◆ GEMINI sample fills in low-luminosity hosts; expectation is that these are low-metallicity as well

NOTE: Does *not* include 118s or 119s; will be addressed






# Fitting Process

5

- ◆ Follows that of Tremonti et al. (2004)
  - ◆ Mask the emission line regions
  - ◆ Find best-fit continuum from large library of models ( $Z$ ,  $T$ , SFH,  $\tau$ ,  $\sigma_{\text{int}}$ )
  - ◆ Subtract continuum; smooth over a sliding window
  - ◆ Fit Gaussians to line regions (Balmer lines; Forbidden lines)






# Fitting Process

6



- ◆ First Cut - S/N (  $>5$  in  $H\alpha$ ;  $H\beta$ ;  $[N II] 6584$ ;  $[O II] 3727$ )
- ◆ Second Cut - BPT diagram (AGN)
  - ◆ Keeping 'composite' for now
- ◆ Assume Case-B recombination for host reddening ( $H\alpha/H\beta$ ) [*EXCEPT* Gemini]
- ◆ Third Cut - 'Global' Z
  - ◆ Fiber size fixed; cutoff used in Kewley & Elston (2008) and Kewley (2005) is 20% g-band
  - ◆ Metallicity Gradients exist in spiral galaxies; lower abundances at higher radii (eg., Henry & Worthey 1999)
  - ◆ Currently tracking, but not cutting (see plot later...)





# Emission Line Measurements <sup>7</sup>


- ◆ Kewley & Dopita (2002):  $[\text{NII}] / [\text{OII}]; R_{23}$
- ◆  $R_{23}$  of M91; KK04 ( $R_{23} = ([\text{OII}] \lambda 3727 + [\text{OIII}] \lambda\lambda 4959, 5007) / \text{H}\beta$ )
- ◆ Depends on models developed using CLOUDY
- ◆ Tremonti (2004) uses simultaneous fits to all emission lines; also CLOUDY





# Results

8



Kewley & Dopita (2002) Diagnostic;  
updated by Kewley & Ellsion (2008)

30 Spec Ia; 6 phot Ia

Trend is detected at  $\sim 2.8\sigma$  for full  
sample;  $\sim 3.5\sigma$  for spec Ia only

DOES NOT include Gemini spectra

DOES NOT include 118s, 119s



star-forming galaxies only



# (less good) Results

9

- ◆ For R23 based methods (KK04; M91),  $1\sigma$  correlation detected ONLY for KK04 spec-only sample
- ◆ Removing the one  $\sim 1/2 Z_{\odot}$  host, this drops to *no* significance.







# R<sub>23</sub> Difficulties

10

I've been able to convert metallicities measured by myself in KD02 system to the Tremonti04 system using KE08 calibration with comparable accuracy to the stated value.

Considerably more scatter - biased - in the conversion from R<sub>23</sub> based methods to KD02





Kewley & Ellison (2008)



# Gemini Spectra

11

- ◆ Observed wavelength range only to  $\sim 6600$  Å; no H $\alpha$ , [N II]
- ◆ Must use R23; cannot determine branch
- ◆ No BPT diagram
- ◆ Reddening from H $\delta$ /H $\beta$
- ◆ Conversion functions of KE08 not valid for *all* R<sub>23</sub> metallicities







# (some) Concerns addressed

12

- ◆ S/N cuts not just a mag cut
- ◆ Possible explanation for  $R_{23}$  conversion difficulties?

- ◆ No inverse correlation between  $Z$  and fiber %





# To Do

13

- ◆ Immediate: Add in GEMINI spectra in some way
- ◆ Almost Immediate: Use Lick indices to get at stellar Z of Ellipticals
- ◆ Merely Urgent: Light Curve properties of SN IIP/IIL; Luminosity functions
- ◆ then ...



...Graduate

14

